

Mathematics 339
April 2010
Professor Cownden
Practice Exam

Please read the following points carefully before proceeding!

1. Please note: Proctors are unable to respond to queries about the interpretation of exam questions. Do your best to answer exam questions as written.
2. You are not allowed any aids other than those included on the last pages of this exam and a Casio 991 calculator or equivalent (gold sticker).
3. Please do not write in red.
4. Print your student number at the top of this page. Do not provide your name.
5. You have 3 hours to complete the exam.
6. Please read all questions carefully before answering them.
7. Answer all questions on the exam paper (both sides); any work not on the exam paper will not be graded. I have tried to ensure that you have enough space on the exam paper to answer each question, but do not feel as if your answer needs to fill out the available space in order to be correct!
8. Marks per question are shown in square brackets after the problem number. The total is 75.
9. Check that your question paper has 9 pages.

Note: This material is copyrighted and is for the sole use of students registered in Math 339 and writing this examination. This material shall not be distributed or disseminated. Failure to abide by these conditions is a breach of copyright and may also constitute a breach of academic integrity under the University Senate's Academic Integrity Policy Statement.

1 The Final Game [15]

Everyone writing the MATH 339 exam has been given this same question. For the purposes of this question assume that there are N students in total writing the exam. There are 80 students enrolled but who knows if they've all shown up.

The game is as follows, each exam writer must select a positive integer (1,2,3...). To win a player must choose the lowest integer n with the property that fewer than n other players have selected the integer n . For example if four people are playing, and one player chooses 1, two players choose 2 and 1 player chooses 3, the player who chooses 1 is the winner and everyone else loses. If on the other hand two players chose 1 and 2 players choose 2 the players who chose 2 would be winners.

1. Find two different Nash Equilibrium for this game, for which all exam writers win.[5]
2. Suppose that $N = 3$. Find all of the mixed strategy Nash equilibria. (Hint use dominated strategies to see which integers should be played with probability zero.)[5]
3. Suppose that every player uses a mixed strategy where they play integer n with probability p_n . (Note that $\sum_{n=1}^N (p_n) = 1$) Write out an expression for the probability of winning in an N player game as a function of the p_n 's.[5]

2 Two-Player Polynomials [20]

Player one must choose a real number, x , between -2 and 2 inclusive. Player two must choose a real number, y , between -3 and 3 inclusive. The payoff to player one is given by

$$\Pi_1(x, y) = x^3y - xy^3 \quad (1)$$

and the payoff to player two is given by

$$\Pi_2(x, y) = x^2y - xy^2 \quad (2)$$

1. Find player one's best response to a given value of y . Hint: This function will be piecewise continuous conditional on the value of y . [5]
2. Find player two's best response to a given value of x . [5]
3. Find all the pairs of values of x and y where both players are playing the best response to each other, aka the intersection of the best response functions, aka the Nash equilibria. [8]
4. What is the best equilibrium for player one, and what is the best equilibrium for player two? [2]

extra workspace

3 ESS and Replicator Equations [40]

Consider the following symmetric two-player matrix game.

	<i>A</i>	<i>B</i>	<i>C</i>
<i>A</i>	(1,1)	(3,4)	(0,2)
<i>B</i>	(4,3)	(0,0)	(0,2)
<i>C</i>	(2,0)	(2,0)	(1,1)

1. Find all the pure strategy Nash equilibria for this game.[2]
2. Under what conditions is a strategy an Evolutionarily Stable Strategy (ESS).[2]
3. Are any of the pure strategies being played at the Nash equilibria you found, Evolutionarily Stable Strategies. Hint: Evolutionary stability is a property of strategies. An equilibrium is only evolutionarily stable if the (mix of) strategies being played there (is) are evolutionarily stable.[4]
4. Let p be the proportion of the population playing A and let q be the proportion of the population playing B , note that the proportion of the population playing C will be $1 - p - q$. Find a mixed strategy Nash equilibria for this game.[2]
5. Is this MSNE an ESS, show your work.[6]
6. Write out the replicator equations for this game. You should have two replicator equations one for the dynamics of p and one for the dynamics of q . [4]
7. Find the equilibria of the system. These are the points where both $\frac{dp}{dt} = 0$ and $\frac{dq}{dt} = 0$. Hint: The mixed strategy is likely an equilibrium point, as are the points where only one strategy type is present in the population.[6]
8. Calculate the Jacobian of the system and use it to determine the stability and type of each of the equilibria of the system.[6]
9. Draw a well **labeled** picture including labeled axis, the easy to draw p and q isoclines, every equilibrium point, and your best guess at what the trajectories look like.[6]
10. Suppose you are in a population that happens to play A with probability $\frac{1}{3}$ B with probability $\frac{1}{3}$, and C with probability $\frac{1}{3}$. What is the best possible strategy that you could adopt for the moment?[2]

(Hint: Since $1 - p - q$ is a proportion we have that $0 \leq 1 - p - q \leq 1$. What constraints does this place on p and q ?, check that your solution is reasonable in light of these constraints.)

extra workspace

more workspace

Classifying Equilibria with a Jacobian

Suppose you have a system of differential equations

$$\begin{aligned}\frac{dp}{dt} &= f(p, q) \\ \frac{dq}{dt} &= g(p, q)\end{aligned}$$

Then the Jacobian of this system is defined to be

$$\mathbf{J} = \begin{pmatrix} \frac{\partial f}{\partial p} & \frac{\partial f}{\partial q} \\ \frac{\partial g}{\partial p} & \frac{\partial g}{\partial q} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

Then let $D = (ad - bc)$ and $T = (a + d)$. If the Jacobian is evaluated at an equilibrium point of the system then using D and T we can classify the type of equilibrium as follows.

1. If $D = 0$ can't say anything about the equilibrium from the Jacobian.
2. If $D < 0$ the equilibrium is a saddle point, and hence unstable. No need to compute T .
3. If $D > 0$ and $T < 0$ the equilibrium is asymptotically stable.
4. If $D > 0$ and $T > 0$ the equilibrium is asymptotically unstable.
5. If $D > 0$ and $T = 0$ the equilibrium point is centre, which is stable but not asymptotically stable. (i.e. not the kind of stable that corresponds to an ESS)
6. If $D > 0$ and $T^2 - 4D < 0$ the equilibrium point is a focus
7. If $D > 0$ and $T^2 - 4D \geq 0$ the equilibrium is a node.

Maximizing a function of two variables

Let $f(p, q)$ be a function of two variables. It's critical points are those which satisfy both of the following equations.

$$\begin{aligned}\frac{\partial f}{\partial p} &= 0 \\ \frac{\partial f}{\partial q} &= 0\end{aligned}$$

To check if a critical point is a local maximum, local minimum, or a saddle point you need to look at the matrix

$$\begin{pmatrix} \frac{\partial^2 f}{\partial p^2} & \frac{\partial^2 f}{\partial p \partial q} \\ \frac{\partial^2 f}{\partial q \partial p} & \frac{\partial^2 f}{\partial q^2} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

Then let $D = ad - bc$ if $D < 0$ the critical point is a saddle point, if $D > 0$ and $\frac{\partial^2 f}{\partial p^2} < 0$ then the critical point is a maximum, if $D > 0$ and $\frac{\partial^2 f}{\partial p^2} > 0$ then the critical point is a minimum.

Binomial distribution

Suppose a coin has a p probability of coming up heads and it is flipped n times. Then the probability of exactly i flips coming up heads is given by $\binom{n}{i}p^i(1-p)^{n-i}$. If your calculator doesn't do this for you automatically $\binom{n}{i} = \frac{n!}{(n-i)!i!}$. This might be helpful in problem one.